

VU Research Portal

Depressive symptoms in an ethnically Diverse cohort of chronic dialysis patients:

Haverkamp, G.L.G.

2017

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Haverkamp, G. L. G. (2017). *Depressive symptoms in an ethnically Diverse cohort of chronic dialysis patients: The role of patient characteristics, cultural and inflammatory factors*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl



Chapter 3

Depressive and anxiety symptoms in Dutch native and immigrant dialysis patients

Wim L. Loosman, Gertrud L.G. Haverkamp, Tessa O. van den Beukel, Tiny Hoekstra, Friedo W. Dekker, Prataap K. Chandie Shaw, Yves F.C. Smets, Louis-Jean Vleming, Pieter M. Ter Wee, Carl E.H. Siegert, Adriaan Honig

Submitted for publication

ABSTRACT

Background

Due to continuing migration there is more interest in the mental health status of immigrants. The aim of this study is to determine the prevalence of depressive/anxiety symptoms in immigrant and native dialysis patients, and to explore if patient characteristics can explain differences.

Methods

For this cross-sectional study patients were selected from the DIVERS study, a prospective cohort study among chronic dialysis patients in the Netherlands. Regions of origin of immigrant patients were defined as African-South American, Asian and European. The Beck Depression Inventory and the Beck Anxiety Inventory were used. Differences between native and immigrant patients were explored using logistic regression models adjusted for patient characteristics.

Results

For 245 native and 249 immigrant patients the prevalence of depressive symptoms was 35 and 50% and of anxiety symptoms 30 and 42%, respectively. Crude ORs for depressive/anxiety symptoms for immigrant patients versus native patients were 1.8 (1.2–2.5) and 1.7 (1.2–2.5), respectively. After adjustment for patient characteristics ORs remained the same.

Conclusion

Clinicians should be aware that immigrant dialysis patients are more prone to develop depressive and anxiety symptoms. Cultural factors might play a role and should therefore be assessed in future research.

INTRODUCTION

Depressive symptoms are highly prevalent in dialysis patients with an estimated occurrence of 39% as was shown by a recent meta-analysis¹. These symptoms are independently associated with mortality², hospitalization³, and impaired health related quality of life (HRQOL)^{4,5}. Depressive symptoms often co-occur with anxiety symptoms⁶. Anxiety symptoms are also highly prevalent in dialysis patients with a prevalence that varies between 13% and 50%⁶⁻¹¹. Anxiety symptoms are also associated with impaired HRQOL⁸ in dialysis patients, and seem to contribute to the relationship between depressive symptoms and impaired HRQOL⁶.

Due to continuing migration there is more interest in the mental and physical health status of immigrants¹². Depressive and anxiety symptoms have been found to vary between native and immigrant groups in a range of settings including the general population and several patients groups¹³⁻¹⁶. In chronic dialysis patients, however, it is unclear whether being immigrant is a risk factor for depressive and anxiety symptoms. A recent meta-analysis¹ showed that the prevalence of depressive symptoms does not seem to differ between white and non-white dialysis patients. However, the studies included in this meta-analysis¹ used ethnic origin but did not include immigrant status and were predominantly US dialysis populations. In these US dialysis populations ethnic origin is often divided in two major groups: African American or Black patients and Caucasian or White patients. In the Netherlands and in many European countries the ethnic and immigrant composition is rather different and diverse, compared to the US dialysis population^{13,17}.

The objective of this study is to determine the prevalence of depressive and anxiety symptoms in native and immigrant dialysis patients in the Netherlands. In addition, we explore if the prevalence of depressive and anxiety symptoms differs between regions of origin of the immigrant patients. Finally, we explore whether differences between dialysis patient groups can be explained by patient characteristics such as demographic and clinical parameters or physical status.

METHODS

Patients

For this study we used cross-sectional data of the "Depression related factors In dialysis patients with Various Ethnicities and Races Study (DIVERS)", which is an observational prospective cohort study among dialysis patients in four urban teaching hospitals and one university hospital in the Netherlands. These hospitals were chosen because of the high percentage of immigrant dialysis patients. Patients were included between June 2012 and December 2013 (prevalent patients) or until October 2014 (incident patients). Patients were included when they were at least 18 years of age, were on dialysis treatment

for at least 90 days, and were able to fill out questionnaires in the Dutch, English, Turkish or Moroccan Arabic language. Patients were excluded if they had cognitive disabilities that interfered with filling out questionnaires. For the current analysis patients were also excluded if they did give permission for examination of their medical records but did not want to fill out questionnaires. The study was approved by the medical ethics committee of the VU university medical centre (approval number: 2010/064).

Patient characteristics

The following demographic and clinical data were obtained from medical records: age, gender, dialysis modality, date of starting dialysis treatment, comorbidity, primary cause of kidney disease, body mass index (BMI), and laboratory measures. Data on smoking status, country of birth, country of birth of both parents and grandparents, marital status, working status and education were collected through a self-report questionnaire. The primary cause of kidney disease was classified according to the codes of the European Renal Association-European Dialysis and Transplantation Association (ERA-EDTA)¹⁸. Co-morbid conditions were scored according to the Davies comorbidity index¹⁹, resulting in a comorbidity index classified into: no, intermediate or severe. Laboratory parameters were collected from routine blood laboratory investigations performed in the individual dialysis centers. To determine patient's physical status we used the physical component summary score (PCS) of the Short Form 12 (SF-12). The SF-12 is a quality of life questionnaire, consisting of a physical and a mental component score. SF-12 is validated among dialysis patients²⁰.

Immigrant status

Patients were classified as native or immigrant based on the country of birth of the patients and of their parents. According to the standard definition of the Statistics Netherlands criteria an individual is considered to be an immigrant if at least one parent was born abroad, regardless of the individual's own country of birth. Patients who are born abroad with at least one parent also born abroad are first generation immigrant patients. Patients who are born in the Netherlands with at least one patient born abroad are second generation immigrants^{21;22}. Patients of whom both parents were born in the Netherlands were classified as native. Moreover, immigrant patients were further classified into region of origin. Patients of whom at least one parent was born in Africa or South America were classified as African-South American immigrants. Patients of whom at least one parent was born in the Middle East or Asia were classified as Asian immigrants. Patients of whom at least one parent was born in Europe, other than the Netherlands, were classified as European immigrants. Surinamese patients of whom at least one grandparent was born in Asia were classified as Asian immigrant. All other Surinamese patients were classified as African-South American immigrant²².

If both parents are born in different countries, the mother's country of birth is used to infer descent.

Depressive and anxiety symptoms

Depressive symptoms were measured using the Beck Depression Inventory (BDI)²³, and symptoms of anxiety were measured using the Beck Anxiety Inventory (BAI)²⁴. Both questionnaires consist of 21 questions which are scored on a 0 – 3 scale. Patients were considered as having depressive and anxiety symptoms when they scored at least 13 points on the BDI²⁵ and 13 points on the BAI^{26,27}, respectively.

Statistical analyses

The three ethnic groups were compared in terms of patient characteristics, symptoms of depression, and anxiety symptoms using the analysis of variance F test or Kruskal-Wallis tests for continuous variables, and a Pearson's chi-square test for categorical variables. Mean with standard deviation or median with interquartile range was used, depending on the underlying distribution. Odds ratios were calculated using logistic regression to determine differences between immigrant groups in the prevalence of depressive and anxiety symptoms, and to determine whether differences between immigrant groups can be explained by demographic variables and clinical characteristics. Multivariable adjustment was done deliberately within the causal pathway in order to explain potential mechanism. We used gradually more extensive multivariable models with demographic and clinical variables, followed by psychosocial variables and the patients physical status.

To obtain complete datasets, missing values for both continuous and categorical variables were imputed with multiple imputation techniques (5 repetitions). With multiple imputation missing data are imputed by a value that is predicted using the patients available characteristics under the condition of missing 'at random'. Missing 'at random' means that the missing value probably depends on other observed patient characteristics. Parathormone, time on dialysis, Kt/V and ferritin were not normally distributed and therefore square root transformed to enter the multiple imputation model. Almost all variables had < 10% missing except for total cholesterol (18%).

A p-value < 0.05 was considered statistically significant. Patient characteristics, depressive and anxiety symptoms were calculated on both the original and imputed dataset. For all statistical analyses SPSS for Windows version 21.0 was used.

Sensitivity analysis

To test the robustness of the results, several sensitivity analyses were performed. Analyses were repeated (1) excluding second generation immigrants (2) excluding

patients who received help filling out the questionnaires, because these patients could give more socially accepted answers; (3) using different cut-off scores (≥ 14 , ≥ 15 and ≥ 16) in our logistic regression models for both the BDI and the BAI; (4) immigrants stratified into Turkish, Moroccan, Surinamese from Asian descent, and Surinamese from African descent because these are the main immigrant group in the Netherlands¹³. Finally, patient characteristics were compared between patients who did and who did not fill out the questionnaires.

RESULTS

Patient characteristics

In total 554 dialysis patients were included the DIVERS study of whom 494 filled out questionnaires and were included in the current analysis (native n=245, immigrant n=249). Patient characteristics of the included native and immigrant patients are presented in table 1. Compared to native patients, immigrants were younger ($p < 0.01$), had a higher prevalence of diabetes ($p < 0.01$), had a lower calcium ($p < 0.01$), and have a different distribution in the cause of end-stage renal disease ($p < 0.01$). The immigrant group consisted of 108 African-South American patients, 114 Asian patients and 27 European patients. Moreover, two hundred and twelve patients (85%) were first generation immigrant patients. The mean length of stay in the Netherlands of these patients was 32 (± 14) years. The countries of origin of all patients are presented in supplementary table 1.

Depressive and anxiety symptoms

The prevalence of depressive and anxiety symptoms varied between immigrant groups (table 2). Thirty-five percent of native patients and 49% of all immigrants had depressive symptoms above threshold (44%, 59% and 33% for African-South American, Asian and European immigrants, respectively). Thirty percent of native patients and 42% of all immigrants had anxiety symptoms above threshold (41%, 47% and 22% for African-South American, Asian and European immigrants, respectively).

Logistic regression analysis immigrants versus natives

The crude OR for scoring above threshold for depressive symptoms using the BDI (≥ 13) for all immigrants versus native dialysis patients was 1.8 (1.2 – 2.5), which implicates that immigrants have a 1.8 higher chance of having depressive symptoms compared to native patients. After stepwise adjustment for demographic characteristics, clinical characteristics, co-morbidity, laboratory values, social characteristics and physical status, ORs remained the same (table 3). After dividing all immigrants in three regions of origin and compared them to natives ORs in the fully adjusted model were 1.6 (0.9 –

2.9), 2.8 (1.6 – 4.9) and 0.9 (0.3 – 2.3) for African-South American, Asian and European immigrants, respectively (table 3).

Table 1. Patient characteristics of 494 native and immigrant dialysis patients

	Native	Immigrant	
Socio demographic	N=245	N=249	P-value
Age, y	69.5 (13.8)	58.9 (15.3)	P < 0.01
Sex, % men	60	61	P = 0.81
Education, % low	41	40	p = 0.72
Married/living together, % yes	56	48	P = 0.06
Having children, % yes	76	79	P = 0.36
Employed, % yes	11	12	P = 0.84
Use of anti-depressants, % yes	9	10	P = 0.56
Clinical			
Incident, % yes	31	29	P = 0.60
Modality, % hemodialysis	89	90	P = 0.76
Time on dialysis, months	17.0 (5.0 – 45.0)	24.0 (6.0 – 65.0)	P = 0.07
BMI, kg/m ²	27 (5.6)	27.2 (6.1)	P = 0.24
Current smoking, % yes	21	17	P = 0.29
Causes of ESRD, %			P < 0.01
Diabetes Mellitus	13	35	
Glomerulonephritis	13	9	
Renal vascular disease	30	25	
Other	45	31	
PCS	36.3 (10.5)	37.8 (10.8)	P = 0.06
Comorbidity			
Davies comorbidity, %			P = 0.06
No	31	23	
Intermediate	51	61	
Severe	19	16	
Diabetes Mellitus, % yes	34	50	P = 0.06
Cardiovascular disease, % yes	29	31	P = 0.15
Laboratory			
Albumin, g/l	36.6 (5.5)	37.5 (5.0)	P = 0.07
Hemoglobin, g/l	7.2 (0.8)	7.1 (0.8)	P = 0.47
PTH, pmol/l	26.9 (14.0 – 45.0)	30.0 (16.0 – 47.7)	P = 0.11
Ca, mmol/l	2.30 (0.14)	2.25 (0.18)	P < 0.01
Phosphorus, mmol/l	1.59 (0.42)	1.63 (0.51)	P = 0.35
Ferritin, ug/l	340 (181 – 562)	365 (173 – 637)	P = 0.51
Cholesterol, mmol/l	4.2 (1.2)	4.1 (1.4)	P = 0.62
Kt/V	1.51 (0.47)	1.52 (0.47)	P = 0.84

Data presented as mean (standard deviation) or median (inter quartile range)

ESRD: End-Stage Renal Disease, BMI: Body Mass Index, PTH: parathyroid hormone,

PCS: Physical Component Summary

Table 2. Depressive and anxiety symptoms in dialysis patient divided in three ethnic groups

<i>Depressive and anxiety symptoms</i>	Native N=245	All immigrants N=249	<u>Region of origin</u> Africa-South America N=108	Asia N=114	Europe N=27
BDI \geq 13 (%)	36	49	44	59	33
BDI total score*	11.2 (7.4)	14.7 (11.0)	13.8 (11.5)	16.5 (11.1)	10.7 (6.3)
BAI \geq 13 (%)	30	42	41	47	22
BAI total score*	9.7 (8.5)	13.3 (11.3)	12.5 (11.3)	15.0 (12.0)	9.2 (6.1)

* Data presented as mean and standard deviation, BDI: beck depression inventory, BAI: beck anxiety

Table 3. Odds ratio's for depressive symptoms (BDI \geq 13) for immigrants versus Native Asian versus native and African-South American versus native dialysis patients

Model	Variables tested	All immigrant versus native	African-South American versus native	Asian versus native	European versus native
1. Unadjusted	ethnicity	1.8 (1.2 – 2.5)	1.4 (0.9 – 2.2)	2.6 (1.6 – 4.1)	0.9 (0.4 – 2.1)
2. Demographic	Model 1 plus age and gender	1.9 (1.3 – 2.8)	1.6 (0.9 – 2.6)	2.8 (1.8 – 4.4)	0.9 (0.4 – 2.2)
3. Clinical	Model 2 plus BMI, dialysis modality, primary kidney disease, smoking, time on dialysis	1.8 (1.2 – 2.7)	1.5 (0.9 – 2.5)	2.6 (1.6 – 4.2)	1.0 (0.4 – 2.3)
4. Co-morbidity	Model 3 plus Davies co-morbidity score	1.8 (1.2 – 2.6)	1.5 (0.9 – 2.5)	2.5 (1.5 – 4.0)	1.0 (0.4 – 2.3)
5. Laboratory	Model 4 plus hemoglobin, calcium, albumin, phosphorus, PTH, ferritin, cholesterol and Kt/V	1.8 (1.2 – 2.8)	1.5 (0.9 – 2.6)	2.5 (1.5 – 4.2)	1.1 (0.5 – 2.7)
6. Social	Model 5 plus marital status, educational level, employment, children	1.8 (1.1 – 2.7)	1.4 (0.8 – 2.5)	2.4 (1.4 – 4.0)	1.2 (0.5 – 2.9)
7. Physical status	Model 6 plus PCS score	1.9 (1.2 – 2.9)	1.6 (0.9 – 2.9)	2.8 (1.6 – 4.9)	0.9 (0.3 – 2.3)

BDI: Beck Anxiety Inventory, BMI: Body Mass Index, PTH: parathyroid hormone,

PCS: Physical Component Summary

The crude OR for scoring above threshold for anxiety symptoms using the BAI (\geq 13) for all immigrants versus native dialysis patients was 1.7 (1.2 – 2.5). After stepwise adjustment for demographic characteristics, clinical characteristics, co-morbidity, laboratory values, social characteristics and physical status, ORs remained the same (table 4). After dividing all immigrants in three regions of origin ORs in the fully adjusted model were 1.9 (1.1 – 3.4), 2.2 (1.3 – 3.9) and 0.7 (0.3 – 1.9) for African-South American, Asian and European immigrants, respectively (table 3).

Table 4. Odds ratio's for anxiety symptoms (BAI ≥ 13) for immigrant versus native Asian versus native and African-South American versus native dialysis patients

Model	Variables tested	All immigrant versus native	African-South American versus native	Asian versus native	European versus native
Unadjusted	ethnicity	1.7 (1.2 – 2.5)	1.6 (>1.0 – 2.6)	2.1 (1.3 – 3.4)	0.7 (0.3 – 1.7)
Demographic	Model 1 plus age and gender	1.8 (1.2 – 2.7)	1.9 (1.1 – 3.1)	2.3 (1.4 – 3.7)	0.7 (0.3 – 1.8)
Clinical	Model 2 plus BMI, dialysis modality, primary kidney disease, smoking, time on dialysis	1.7 (1.2 – 2.7)	1.8 (1.1 – 3.1)	2.2 (1.4 – 3.7)	0.7 (0.3 – 1.8)
Co-morbidity	Model 3 plus Davies co-morbidity score	1.7 (1.1 – 2.6)	1.8 (>1.0 – 3.0)	2.1 (1.3 – 3.5)	0.7 (0.3 – 1.7)
Laboratory	Model 4 plus hemoglobin, calcium, albumin, phosphorus, PTH, ferritin, cholesterol and Kt/V	1.8 (1.2 – 2.8)	1.9 (1.1 – 3.3)	2.2 (1.3 – 3.7)	0.8 (0.3 – 2.2)
Social	Model 5 plus marital status, educational level, employment, children	1.7 (1.1 – 2.6)	1.7 (0.9 – 3.0)	2.0 (1.2 – 3.5)	0.8 (0.3 – 2.3)
Physical status	Model 6 plus PCS score	1.7 (1.1 – 2.7)	1.9 (1.1 – 3.4)	2.2 (1.3 – 3.9)	0.7 (0.2 – 1.9)

BAI: Beck Anxiety Inventory, BMI: Body Mass Index, PTH: parathyroid hormone,
PCS: Physical Component Summary

Sensitivity analysis

By excluding second generation immigrants differences between native and immigrant patients remained the same (data not shown). Twenty five percent of the patients needed help filling out the questionnaires. By excluding these patients differences between native and immigrant patients also remained the same (data not shown). After using different cut-off scores for BDI and BAI (≥ 14 , ≥ 15 and ≥ 16), differences between groups became more pronounced when a higher cut-off for BDI or BAI was used. Furthermore, after selecting immigrants from Turkey, Morocco, Surinam from Asian descent and Surinam from African descent the prevalence of depressive symptoms was 56, 77, 70 and 44%, respectively and for anxiety symptoms 44, 71, 48 and 44%, respectively. Finally, no differences for patient characteristics were found between the 494 patients who did and the 60 patients who did not fill out the questionnaires (data not shown).

DISCUSSION

This observational study examined the prevalence of depressive and anxiety symptoms in native and immigrant chronic dialysis patients. Compared to native patients, immigrant patients were almost two times more likely to report depressive and anxiety symptoms. These differences were more pronounced in Asian patients followed by African-South American patients. No differences in the prevalence of depressive and anxiety symptoms were found between European immigrants and native patients. We showed that patients characteristics do not appear to explain the differences between groups.

Our cohort shows that immigrant dialysis patients have a higher prevalence of depressive and anxiety symptoms compared to native dialysis patients. A recent meta-analysis¹ showed no difference between white and non-white dialysis patients. However, in these cohorts ethnicity is reported instead of immigrant status which is not necessarily the same. Moreover, in the US general population, a lower prevalence of depressive symptoms was shown in immigrants compared to natives²⁸. This contradiction could be explained by the differences in the ethnic and immigrant composition of many European countries compared to the US^{12,16}. Conclusively, immigrant status seems to be a risk factor for developing depressive and anxiety symptoms in Dutch chronic dialysis patients.

Immigrant dialysis patients in our cohort were divided in three groups according to region of origin. The first group was from Africa-South America. African-South American immigrants dialysis patients have a slightly higher, but statistically not significant, prevalence of depressive symptoms than native patients. The second region of origin in our cohort was Asia. The Asian immigrant dialysis patients in comparison to native patients had the highest prevalence of depressive symptoms of all immigrant groups. To the best of our knowledge there are no studies who reported prevalence rates of depressive symptoms in Asian immigrant dialysis patients. Both African-South American and Asian patients have a higher prevalence of anxiety symptoms in our cohort. In the few studies that describe the prevalence of anxiety symptoms in dialysis patients differences between ethnic groups could not be made because of the small sample of different ethnic groups⁶⁻¹¹. The third region of origin in our cohort was Europe. European immigrants does not seem to differ from native patients, but the sample size is small. In sum, in addition to immigrant status the region of origin seems to be important for assessing the risk of mental health problems in dialysis patients.

Socio-demographic and clinical patient characteristics could not explain the differences between immigrant and native patients as was shown by a series of logistic regression models. There could be several explanations for the difference between native and immigrant patients. First, perceived ethnic discrimination (PED) may be an explanation, because PED contributes considerably to depression in ethnic minority groups in the Netherlands. Second, it has been shown that less skills for living in the Dutch society, and more feelings of loss are associated with the presence of both depressive and anxiety

symptoms in immigrant chronic dialysis patients^{29,30}. Finally, religion might play a role in the difference between immigrant and native patients³¹⁻³³. Ramirez et al³² showed that in dialysis patients religious struggle was independently associated with greater psychological distress and impaired HRQOL. For clinical research it is important to assess immigrant status besides ethnic origin, and in clinical practice it is important for nephrologists to be aware that immigrant patients are more prone to develop depressive and anxiety symptoms.

A few limitations should be noted for this study. First, the broad spectrum of countries used to define groups makes it difficult to generalize these results for specific immigrant groups. Generalisation is especially difficult for Turkish and Moroccan immigrants which are the main immigrant groups in the Netherlands. Second, the presentation of a depression and anxiety disorder could differ across cultures. However, the BDI was validated in a comparable multi-ethnic cohort of dialysis patients²⁵. Third, some participants filled out the questionnaire with help which makes the risk that participants give socially accepted answers higher. However, differences between ethnic groups remained the same after excluding the patients who received help with filling out the questionnaire.

In conclusion, immigrant dialysis patients have a higher prevalence of depressive and anxiety symptoms than native dialysis patients. Differences between native and immigrant patients were more pronounced in Asian patients. Clinicians should be aware that immigrant dialysis patients are more prone to develop depressive and anxiety symptoms. Patients characteristics could not explain the differences between immigrant and native dialysis patients. Cultural factors might play a role and should therefore be assessed in future research.

REFERENCES

1. Palmer S, Vecchio M, Craig JC, Tonelli M, Johnson DW, Nicolucci A, Pellegrini F, Saglimbene V, Logroscino G, Fishbane S, Strippoli GF. Prevalence of depression in chronic kidney disease: systematic review and meta-analysis of observational studies. *Kidney Int* 2013; 84(1):179-191.
2. Farrokhi F, Abedi N, Beyene J, Kurdyak P, Jassal SV. Association between depression and mortality in patients receiving long-term dialysis: a systematic review and meta-analysis. *Am J Kidney Dis* 2014; 63(4):623-635.
3. Lacson E Jr, Bruce L, Li NC, Mooney A, Maddux FW. Depressive affect and hospitalization risk in incident hemodialysis patients. *Clin J Am Soc Nephrol* 2014; 9(10):1713-1719.
4. Garcia-Llana H, Remor E, Del PG, Selgas R. The role of depression, anxiety, stress and adherence to treatment in dialysis patients health-related quality of life: a systematic review of the literature. *Nefrologia* 2014; 34(5):637-657.
5. Cruz LN, Fleck MP, Polanczyk CA. Depression as a determinant of quality of life in patients with chronic disease: data from Brazil. *Soc Psychiatry Psychiatr Epidemiol* 2010; 45(10):953-961.
6. Preljevic VT, Osthus TB, Os I, Sandvik L, Opjordsmoen S, Nordhus IH, Dammen T. Anxiety and depressive disorders in dialysis patients: association to health-related quality of life and mortality. *Gen Hosp Psychiatry* 2013; 35(6):619-624.
7. Kutner NG, Fair PL, Kutner MH. Assessing depression and anxiety in chronic dialysis patients. *J Psychosom Res* 1985; 29(1):23-31.
8. Cukor D, Coplan J, Brown C, Friedman S, Newville H, Safier M, Spielman LA, Peterson RA, Kimmel PL. Anxiety disorders in adults treated by hemodialysis: a single-center study. *Am J Kidney Dis* 2008; 52(1):128-136.
9. Bossola M, Ciciarelli C, Di SE, Conte GL, Vulpio C, Luciani G, Tazza L. Correlates of symptoms of depression and anxiety in chronic hemodialysis patients. *Gen Hosp Psychiatry* 2010; 32(2):125-131.
10. Zhang M, Kim JC, Li Y, Shapiro BB, Porszasz J, Bross R, Feroze U, Upreti R, Martin D, Kalantar-Zadeh K, Kopple JD. Relation between anxiety, depression, and physical activity and performance in maintenance hemodialysis patients. *J Ren Nutr* 2014; 24(4):252-260.
11. Feroze U, Martin D, Kalantar-Zadeh K, Kim JC, Reina-Patton A, Kopple JD. Anxiety and depression in maintenance dialysis patients: preliminary data of a cross-sectional study and brief literature review. *J Ren Nutr* 2012; 22(1):207-210.
12. Vaughn LM, Jacquez F, Lindquist-Grantz R, Parsons A, Melink K. Immigrants as Research Partners: A Review of Immigrants in Community-Based Participatory Research (CBPR). *J Immigr Minor Health* 2016.
13. de Wit MA, Tuinebreijer WC, Dekker J, Beekman AJ, Gorissen WH, Schrier AC, Penninx BW, Komproe IH, Verhoeff AP. Depressive and anxiety disorders in different ethnic groups: a population based study among native Dutch, and Turkish, Moroccan and Surinamese migrants in Amsterdam. *Soc Psychiatry Psychiatr Epidemiol* 2008; 43(11):905-912.
14. Missinne S, Bracke P. Depressive symptoms among immigrants and ethnic minorities: a population based study in 23 European countries. *Soc Psychiatry Psychiatr Epidemiol* 2012; 47(1):97-109.
15. Haverkamp GL, Torensma B, Vergouwen AC, Honig A. Psychological Distress in the Hospital Setting: A Comparison between Native Dutch and Immigrant Patients. *PLoS One* 2015; 10(6):e0130961.
16. Tarricone I, Stivanello E, Poggi F, Castorini V, Marseglia MV, Fantini MP, Berardi D. Ethnic variation in the prevalence of depression and anxiety in primary care: a systematic review and meta-analysis. *Psychiatry Res* 2012; 195(3):91-106.

17. Goodkin DA, Bragg-Gresham JL, Koenig KG, Wolfe RA, Akiba T, Andreucci VE, Saito A, Rayner HC, Kurokawa K, Port FK, Held PJ, Young EW. Association of comorbid conditions and mortality in hemodialysis patients in Europe, Japan, and the United States: the Dialysis Outcomes and Practice Patterns Study (DOPPS). *J Am Soc Nephrol* 2003; 14(12):3270-3277.
18. van Dijk PC, Jager KJ, de CF, Collart F, Cornet R, Dekker FW, Gronhagen-Riska C, Kramar R, Leivestad T, Simpson K, Briggs JD. Renal replacement therapy in Europe: the results of a collaborative effort by the ERA-EDTA registry and six national or regional registries. *Nephrol Dial Transplant* 2001; 16(6):1120-1129.
19. Davies SJ, Russell L, Bryan J, Phillips L, Russell GI. Comorbidity, urea kinetics, and appetite in continuous ambulatory peritoneal dialysis patients: their interrelationship and prediction of survival. *Am J Kidney Dis* 1995; 26(2):353-361.
20. Loosman WL, Hoekstra T, van Dijk S, Terwee CB, Honig A, Siegert CE, Dekker FW. Short-Form 12 or Short-Form 36 to measure quality-of-life changes in dialysis patients? *Nephrol Dial Transplant* 2015.
21. Statistics Netherlands. Someone with a foreign background. Available from: URL:<http://www.cbs.nl/en-GB/menu/methoden/begripen/default.htm?Languageswitch=on&ConceptID=37>. Accessed 2015 May.
22. Stronks K, Kulu-Glasgow I, Agyemang C. The utility of 'country of birth' for the classification of ethnic groups in health research: the Dutch experience. *Ethn Health* 2009; 14(3):255-269.
23. Beck AT, Ward CH, Mendelson M, Mok J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry* 1961; 4:561-571.
24. Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychol* 1988; 56(6):893-897.
25. Loosman WL, Siegert CE, Korzec A. Validity of the Hospital Anxiety and Depression Scale and the Beck Depression Inventory for use in end-stage renal disease patients. *Br J Clin Psychol* 2010; 49:507-516.
26. Wang MY, Chan SF, Chang LI, Chen TH, Tsai PS. Better sleep quality in chronic haemodialyzed patients is associated with morning-shift dialysis: a cross-sectional observational study. *Int J Nurs Stud* 2013; 50(11):1468-1473.
27. Che HH, Lu ML, Chen HC, Chang S.W., Lee YJ. Validation of the Chinese version of the Beck Anxiety Inventory. *Formos J Med* 2006; 10:447-454.
28. Wong EC, Miles JN. Prevalence and correlates of depression among new U.S. immigrants. *J Immigr Minor Health* 2014; 16(3):422-428.
29. Koneru VK, de Mamani AGW, Flynn PM, Betancourt H. Acculturation and mental health: current findings and recommendations for future research. *Appl Prev Psychol* 2007; 12:76-96.
30. Haverkamp GL, Loosman WL, van den Beukel TO, Hoekstra T, Dekker FW, Chandie Shaw P.K., Smets Y. The association of acculturation and depressive and anxiety symptoms in immigrant chronic dialysis patients. *Gen Hosp Psychiatry* 2015.
31. Ramirez SP, Macedo DS, Sales PM, Figueiredo SM, Daher EF, Araujo SM, Pargament KI, Hyphantis TN, Carvalho AF. The relationship between religious coping, psychological distress and quality of life in hemodialysis patients. *J Psychosom Res* 2012; 72(2):129-135.
32. Patel SS, Shah VS, Peterson RA, Kimmel PL. Psychosocial variables, quality of life, and religious beliefs in ESRD patients treated with hemodialysis. *Am J Kidney Dis* 2002; 40(5):1013-1022.
33. Fernandez A, Loukas A. Acculturation and religious coping as moderators of the association between discrimination and depressive symptoms among Mexican-American vocational students. *J Immigr Minor Health* 2014; 16(6):1290-1293.

SUPPLEMENTARY MATERIAL

Supplementary table 1. Countries of origin of at least one parent of 494 dialysis patients included in the DIVERS study.

Native N = 245	Africa -South America N = 108	Asia N = 114	Europe N = 27
245 The Netherlands	52 Surinam*	46 Surinam*	14 Germany
	17 Morocco	32 Indonesia	3 Belgium
	12 Netherlands Antilles	16 Turkey	3 Poland
	5 Ghana	6 Pakistan	2 Portugal
	5 Somalië	5 India	1 Scotland
	3 Egypte	2 Afghanistan	1 Great Britain
	2 Colombia	2 Filippijnen	1 Slovakia
	2 Algerije	1 China	1 Romania
	2 Nigeria	1 Nepal	1 Italy
	2 Soedan	1 Armenia	
	1 Guinee	1 Iran	
	1 Eritrea	1 Irak	
	1 Liberia		
	1 Uruguay		
	1 Brasil		
	1 Sierra Leone		

* Surinamese patients of whom at least one grandparent was born in Asia were classified as Asian. All other Surinamese patients were classified as African-South American.

